



**Supports absorption
of dietary
calcium/phosphorus**

**Maintain bird
welfare**

**Aids vitamin D
metabolism**

**Gut health
support**

D-Plus

A complementary feedstuff containing vitamins and 25-hydroxy vitamin D3 for use in all poultry. Designed for strategic use where supply of nutrients from the feed is inadequate for optimal bone development, shell quality, good health and performance.

**Vitamin D
25 (OH) D3
Vitamin B
Vitamin K
Vitamin C
Medium Chain Fatty Acids**

Objectives:

- Development of strong bones and shell.
- Support absorption of dietary calcium and phosphorus.
- Reduce financial loss due to carcass/egg downgrading.
- Maintain bird welfare.

Product Indications:

- High incidence of leg culls or factory rejects.
- Reduced movement.
- Poor hatchability.
- Shell cracks or reduced quality.

Technical Interest:

Strong bones: Fast growing birds require adequate support to maintain feeding, active behaviour, health and welfare. A strong bone structure, particularly legs can increase mobility resulting in positive feeding along with welfare affects, supporting performance.

Shell weakness: Point of lay, optimum lay, and later lay persistence rely on ensuring correct balance of calcium and phosphorus. Adequate absorption and metabolism must be maintained to help minimise hairline cracks, broken shells or rejects. Provision of good shell quality also assists hatchability of chicks.



For Poultry

A complementary feedstuff containing vitamins and 25-hydroxy vitamin D3 for use in all poultry. Designed for strategic use where supply of nutrients from the feed is inadequate for optimal bone development, shell quality, good health and performance.

Benefits:

- Contributes to utilisation of calcium and phosphorus from the feed.
- Contributes to calcification of bones and egg shells.

D-Plus

When to apply:

✓
2 days during fast growth period
(Broilers)

✓
Strategically Day 11-14 and/or 21-24
(Broilers)

✓
2 days during welfare challenges associated with poor mobility
(Broilers & Layers)

✓
2 days per week for 5 weeks during peak lay
(Layers)

✓
2 days per week when poor shell quality is observed
(Layers)

D-Plus



Application:

Add 1,000ml of **D-Plus** liquid to 1,000 litres of clean drinking water. Ensure Dosatron™ stock solution is calibrated for appropriate uptake. No manual stirring is required. Product can be used as standard management practice or during individual cycle cases of digestive stress according to the discretion of the farm manager. To maintain drinking lines and water quality, any water application should be followed with a suitable 1:50,000 dilution of sanitiser.

Storage:

Store in cool and dry place away from direct light. Keep away from children.

Shelf Life: 24 months from date of manufacturing.

Pack Size: 1L Bottle.