



---

**Aid digestive  
function**

---

---

**Support energy  
levels and flock  
uniformity**

---

---

**Assist normal  
liver function**

---

## Nutaliver

A complementary liquid feedstuff for all poultry to provide energy, help support liver function and aid digestive activity during periods of poor performance, times of stress or reduced feed intake.

Essential oils  
Plant extracts  
Sorbitol  
Choline  
Methione  
Betaine

### Objectives:

- Support performance during digestive stress.
- Assist normal liver function.
- Help support healthy gut function.

### Product Indications:

- Help encourage healthy digestion during stress situations.
- Support metabolism by providing additional energy.
- Support inactive or poor performing birds.

### Technical Interest:

**Digestive function:** Ingredients which help gut transit, support of gut enzymes and secretion of bile salts into the digestive tract allow for better processing of nutrients, including fats or oils.

**Inactivity and uniformity:** Supply of energy as alternatives to glucose or fat while supplying energy to support FCR.

**Liver support:** Energy supplies independent to the liver reduce the overall liver load. Supporting liver health can allow effective metabolism of fats and assist with detoxification, particularly in layers.



For Poultry

A complementary liquid feedstuff for all poultry to provide energy, help support liver function and aid digestive activity during periods of poor performance, times of stress or reduced feed intake.

#### Benefits:

- Relieve FCR impact associated with poor feed utilisation.
- Aid egg quality and number by supporting liver function.
- Help maintain active feeding behaviour.
- No withdrawal period.

## Nutaliver

When to apply:

✓  
**3-5 days**  
during early  
growth period  
(Broilers)

✓  
**2-3 days**  
during low  
growth or  
poor FCR  
(Broilers)

✓  
**3 days**  
during digestive  
challenge  
(Broilers & Layers)

✓  
**2 days**  
per week  
for 5 weeks  
to point of lay  
(Layers)

✓  
**2 days per**  
**week**  
during poor liver  
function  
(Broilers & Layers)



#### Application:

Add 1000-2000ml of **Nutaliver** liquid to 1000lt of clean drinking water. Ensure Dosatron™ stock solution is calibrated for appropriate uptake. No manual stirring is required. Product can be used as standard management practice or during individual cycle cases of digestive stress according to farm managers discretion. To maintain drinking lines and water quality, any water application should be followed with a suitable 1:50,000 dilution of sanitiser.

#### Storage:

Store in cool and dry place away from direct light. Keep away from children.

**Shelf Life:** 24 months from date of manufacturing.

**Pack Size:** 5L Jerrycan.

Nutaliver